

# Anaphylaxis Emergency Plan

(All sections must be completed.)

Student Name: \_\_\_\_\_



**This person has a potentially life-threatening allergy (anaphylaxis) to:**

(Check the appropriate boxes)

Food(s): \_\_\_\_\_  
\_\_\_\_\_

Insect stings

Other: \_\_\_\_\_

**Epinephrine Auto-Injector:** Expiry Date: \_\_\_\_\_/\_\_\_\_\_/\_\_\_\_\_

**Dosage:**  EpiPen Jr. 0.15 mg     EpiPen 0.30 mg

**Location of Auto-Injector(s):** \_\_\_\_\_

**Previous anaphylactic reaction:** Person is at greater risk.

**Asthmatic:** Person is at greater risk. If person is having a reaction and has difficulty breathing, give epinephrine auto-injector before asthma medication.

**A person having an anaphylactic reaction might have ANY of these signs and symptoms:**

- **Skin system:** hives, swelling (face, lips, tongue), itching, warmth, redness
- **Respiratory system (breathing):** coughing, wheezing, shortness of breath, chest pain or tightness, throat tightness, hoarse voice, nasal congestion or hay fever-like symptoms (runny, itchy nose and watery eyes, sneezing), trouble swallowing
- **Gastrointestinal system (stomach):** nausea, pain or cramps, vomiting, diarrhea
- **Cardiovascular system (heart):** paler than normal skin colour/blue colour, weak pulse, passing out, dizziness or lightheadedness, shock
- **Other:** anxiety, sense of doom (the feeling that something bad is about to happen), headache, uterine cramps, metallic taste

*Early recognition of symptoms and immediate treatment could save a person's life.*

**Act quickly. The first signs of a reaction can be mild, but symptoms can get worse very quickly.**

1. **Give epinephrine auto-injector** (e.g. EpiPen) at the first sign of a known or suspected anaphylactic reaction (see attached instruction sheet).
2. **Call 9-1-1** or local emergency medical services. Tell them someone is having a life-threatening allergic reaction.
3. **Give a second dose of epinephrine** as early as 5 minutes after the first dose if there is no improvement in symptoms.
4. **Go to the nearest hospital immediately (ideally by ambulance)**, even if symptoms are mild or have stopped. The reaction could worsen or come back, even after proper treatment. Stay in the hospital for an appropriate period of observation as decided by the emergency department physician (generally about 4-6 hours).
5. **Call emergency contact person (e.g. parent, guardian).**

**Emergency Contact Information**

Name	Relationship	Home Phone	Work Phone	Cell Phone

The undersigned patient, parent or guardian authorizes any adult to administer epinephrine to the above-named student in the event of an anaphylactic reaction as described above. This protocol has been recommended by the patient's physician.

\_\_\_\_\_  
Patient/Parent/Guardian Signature

\_\_\_\_\_  
Physician Signature  On file

\_\_\_\_\_  
Date

\_\_\_\_\_  
Date